

Original Article

# Magnetic Resonance Imaging Compared with Rectal Endoscopic Sonography for the Prediction of Infiltration Depth in Colorectal Endometriosis

Arane Kim, MD, Pedro Fernandez, MD, Brigitte Martin, MD, Laurent Palazzo, MD, Lara Ribeiro-Parenti, MD, Francine Walker, MD, Margot Bucau, MD, Helene Collinot, MD, Dominique Luton, MD, PhD, and Martin Koskas, MD, PhD

From the Department of Obstetrics and Gynecology (Drs. Kim, Collinot, Luton, and Koskas), Departments of Radiology (Dr. Fernandez), Digestive Surgery (Dr. Ribeiro-Parenti), Pathology, Bichat Hospital, Paris, France (Drs. Walker and Bucau), Institut de Radiologie de Paris, Paris, France (Dr. Martin), Clinique du Trocadéro, Paris, France (Dr. Palazzo), and Paris Diderot University, Paris, France (Dr. Koskas).

**ABSTRACT** **Study Objective:** To compare the accuracies of magnetic resonance imaging (MRI) and rectal endoscopic sonography (RES) in the prediction of the infiltration depth of colorectal endometriosis.

**Design:** A retrospective cohort study (Canadian Task Force classification II-2).

**Setting:** A university teaching hospital.

**Patients:** Forty patients with symptomatic deep infiltrating endometriosis (DIE) of the rectum who underwent colorectal resection were included.

**Interventions:** All patients underwent abdominopelvic MRI and RES preoperatively to assess the infiltration depth of colorectal endometriosis, and segmental resection of the rectosigmoid by laparoscopy was performed if RES showed bowel invasion. Sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), positive and negative likelihood ratios (LRs), and intermethod agreement were calculated for DIE muscularis and submucosal/mucosal infiltration confirmed by histopathological analysis.

**Measurements and Main Results:** For MRI detection of DIE muscularis infiltration, the sensitivity, specificity, PPV, NPV, and negative LR were 68%, 100%, 100%, 20%, and 0.32, respectively. For the MRI detection of DIE submucosal/mucosal involvement, the sensitivity, specificity, PPV, NPV, and positive and negative LR were 47%, 81%, 69%, 63%, 2.49, and 0.65, respectively. The PPV of RES detection of DIE muscularis infiltration was 93%. For the RES detection of DIE submucosal/mucosal layers, the sensitivity, specificity, PPV, NPV, and positive and negative LR were 79%, 48%, 58%, 71%, 1.51, and 0.44, respectively.

**Conclusion:** In the current study, MRI is valuable for detecting endometriosis of the rectum but is less accurate in detecting submucosal/mucosal involvement than RES. Magnetic resonance imaging was not successful for preoperative determination of segmental resection versus a more conservative approach. When bowel involvement is detected by MRI, RES is not essential. When symptoms suggest DIE in patients without intestinal lesions detected by MRI, RES is necessary to exclude bowel invasion. Journal of Minimally Invasive Gynecology (2017) 24, 1218–1226 © 2017 AAGL. All rights reserved.

**Keywords:** Deep infiltrating endometriosis; Imaging modality; Laparoscopic colorectal resection

Intestinal endometriosis concerns 5% to 12% of patients with endometriosis [1] and is defined as endometriosis involving the bowel only if the muscularis layer is invaded. In

The authors declare that they have no conflict of interest.

Corresponding author: Arane Kim, MD, Department of Obstetrics and Gynecology, Bichat Hospital, CHU Bichat Claude Bernard, 46 rue Henri-Huchard, 75018 Paris, France.

E-mail: [aranekim@gmail.com](mailto:aranekim@gmail.com)

Submitted May 27, 2017. Accepted for publication July 13, 2017.

Available at [www.sciencedirect.com](http://www.sciencedirect.com) and [www.jmig.org](http://www.jmig.org)

1553-4650/\$ — see front matter © 2017 AAGL. All rights reserved.

<https://doi.org/10.1016/j.jmig.2017.07.026>

such patients, dyschezia, rectal bleeding, cyclic defecation pain, constipation, and/or diarrhea are typical [2].

In patients with intestinal endometriosis, segmental rectal resection with colorectal anastomosis reduces recurrence and improves digestive and gynecologic symptoms and quality of life [3]. However, it is associated with complications such as rectovaginal fistulae (1.8%–2.7%), anastomotic leakages (1.5%–1.9%), and pelvic abscesses (0.34%–1%) [4,5]. Alternatively, rectal shaving results in less morbidity while preserving organs, nerves, and vascular blood supply [6,7].

Transvaginal sonography, rectal endoscopic sonography (RES), and magnetic resonance imaging (MRI) are used to detect and localize intestinal endometriosis [8,9]. Transvaginal sonography is the first-line imaging modality for endometriosis assessment, whereas MRI and RES are second-line for the detection of upper digestive lesions and the depth of colorectal lesions, respectively [9,10].

Although RES is a beneficial diagnostic tool for colorectal endometriosis, it has been suggested that MRI is the best non-invasive method for evaluating locations of pelvic endometriosis [11]. However, few studies have evaluated the accuracy of imaging to predict the infiltration depth of colorectal endometriosis [10,12–15]. In the present study, we aimed to compare the accuracies of MRI and RES in predicting the depth of colorectal endometriosis. This information could be useful to identify patients who do not require radical segmental rectal resection but rather rectal shaving only or a transmural local excision with primary closure.

## Methods

This retrospective, single-center study was conducted between 2012 and 2015 and included 40 consecutive patients who underwent colorectal resection for deep infiltrating endometriosis (DIE). All patients with symptoms of DIE (eg, dysmenorrhea, dyspareunia, chronic pelvic pain, and dysuria) and colorectal endometriosis (eg, dyschezia and cyclical rectal bleeding) underwent a physical examination and preoperative abdominopelvic MRI and RES.

Histologic examination was used as validation of bowel endometriosis and infiltration depth. Only patients with DIE muscularis involvement noted on preoperative RES underwent a segmental resection of the rectosigmoid performed by laparoscopy. The study was approved by the local institutional review board, and all patients were informed that their deidentified data would be collected for research purposes.


## RES Protocol

RES was performed by a single examiner (L.P.) with extensive experience in RES for DIE for all patients with symptoms possibly related to digestive rectosigmoid endometriosis. The sonographer was blinded to the results of MRI.

The echoendoscope used was a flexible Olympus (Rungis, France) radial electronic with a 14.5-mm-diameter end, allowing 360° cuts on a perpendicular plane and oblique anterior echoendoscopic visibility [12]. The device was coupled to an Aloka Alpha10 console (Aloka, Saint-Priest, France). The standard probe frequency used to detect nodules was 6 MHz, and 10 MHz was used when refining of an interpretation was needed (eg, to visualize the mucosa or submucosa). A Normacol (Norgine, Rueil Malmaison, France) (sodium dihydrogen phosphate) enema was performed 2 hours before the examination to reduce gas and fecal material-related artifacts. Ultrasonography was performed without sedation. However, in the event of significant patient anxiety or fore-

**Fig. 1**

RES of the invasion of the muscularis layer by an endometriotic nodule of the rectosigmoid.




seeable difficulties performing the test, sedation with propofol was available for fasting patients [12].

Normal anatomy appears as follows on ultrasound for the rectosigmoid [12]: the interface between the serosa and muscular layer appears as an external hyperechoic line (thinner than 0.5 mm), hypoechoic external and internal smooth muscle layers (2 mm) are separated by a hyperechoic line, and hyperechoic mucosal and submucosal layers are separated by hypoechoic muscularis mucosa. Digestive parietal invasion was defined by the presence of a hypoechoic nodule colonizing the rectal or sigmoid wall associated with a thickening of the muscular layer. Mucosal or submucosal infiltrations were characterized by interruption of their hyperechoic line [12] (Figs. 1 and 2).

**Fig. 2**

RES of the invasion of the mucosa layer by an endometriotic nodule of the anterior rectum.



### MRI Protocol

MRI was reinterpreted by a single radiologist blind to both RES and histologic results. We aimed to find out if MRI was more accurate than RES or not when RES was positive.

MRI examination was performed with a 1.5-Tesla MRI device with 2- to 5-mm-thick sections and a 1- to 3-mm gap. Intrarectal or intravaginal gel-based preparation was used, although not in all subjects, and antispasmodic drugs (Glucagen; Novo Nordisk, Paris, France) were occasionally injected to reduce peristalsis.


The MRI protocol included a series of at least 3 spatial imaging planes with acquisitions including T2-weighted and T1-weighted images with and without fat suppression. Gadolinium injection was not administered for every scan (Figs. 3 and 4).

The main aspects of MRI of endometriosis were described as hyperintense foci corresponding to hemorrhagic signs on T1-weighted and fat-suppressed T1-weighted MRI, hyperintense cavities on T2-weighted MRI, hypointense signs with low enhancement on T1-weighted and T2-weighted MRI with retractable or spiculated signs, and fibrosis with signal intensity close to pelvic muscle on T1- and T2-weighted MRI [16].

Endometriosis lesions of the anterior wall of the rectosigmoid colon were observed as disappearance of the hypointense signal on T2-weighted images. The presence of nodules extending on the anterior and inferior wall of the rectosigmoid colon showing contrast enhancement on T1-weighted MRI was also observed. Muscular or mucosal and submucosal infiltrations were characterized by the exten-

**Fig. 3**

MRI of the sagittal sequence T2-weighted, retractable nodule of the rectum extending on the submucosa with a T2 hypointense signal. Double arrows = retractable nodule of the rectum.



**Fig. 4**

MRI of the oblique coronal T2-weighted sequence. Rectal nodule (arrow). Mucosa-extended rectal wall lesion (double arrow).



sive depth of muscular thickening of the rectal wall (isosignal). All patients underwent laparoscopy with segmental resection of the rectosigmoid by the same surgical team using similar operative techniques. Antibiotic prophylaxis with cefazolin 2 g was administered, and a 10-mm laparoscope was used in the umbilical position with 3 other trocars. Segmental resection was guided by lesions in the digestive tract diagnosed by preoperative imaging and those that were macroscopically identified intraoperatively. A minilaparotomy was performed to make the anastomosis using a circular mechanical clamp.


Histologic examinations of the resected bowel were performed by the same experienced pathologist (F.W.) with extensive experience in recognizing DIE who was blinded to the results of MRI and RES. Endometriosis was defined by the presence of fibrosis and muscular hyperplasia in association with ectopic endometrial tissue (ie, glandular and stromal structures on light microscopy) as shown in Figs. 5 and 6. The results of RES, MRI, and pathologic analysis of the excised tissues were compared. The maximum depth reached by endometriosis in the bowel was considered.

### Statistical Analysis

The sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV), test accuracy, positive and negative likelihood ratios (LRs), and corresponding 95% confidence interval (CI) of MRI and RES were reported for each location of endometriosis and invasion depth (muscular layers and the colorectal submucosal/mucosal layers).

**Fig. 5**

Submucosa (left side \*), muscular (middle \*), and subserosa (right side \*) infiltration of deep infiltrating endometriosis of the rectum. Zoom  $\times 6$ .




Because only patients with muscular lesions on RES were included, it was not relevant to calculate the sensitivity, specificity, NPV, and LRs for the muscularis layer.

Intermethod agreement was calculated with the Cohen kappa coefficient. The degree of agreement was defined according to Landis and Koch [17] (<0, no agreement; 0.00–0.20, slight agreement; 0.21–0.40, fair agreement; 0.41–0.60, moderate agreement; 0.61–0.80, substantial agreement; and 0.81–1.00, excellent agreement).

**Fig. 6**

Submucosa infiltration of deep infiltrating endometriosis of the rectum (\*). Zoom  $\times 50$ .



Agreement between RES and MRI was calculated using 3 categories: no bowel wall invasion, muscularis, and submucosal/mucosal. Agreement between the radiologic examination (RES or MRI) and histologic results using the same categories was also calculated.

**Results**

Forty patients with a median age of 33 years were analyzed. The most common symptoms were dysmenorrhea (80%), dyschezia and chronic pelvic pain (both 70%), and dyspareunia (68%) (Table 1). All patients had a maximal interval of 4 months between imaging (MRI and RES) and surgery.

Histopathological examination showed that 18 (45%) of the 40 patients had muscular invasion only, and 19 (48%) had associated submucosal or mucosal infiltration. In 3 patients, serosal involvement only was observed and no muscularis invasion (Tables 2 and 3).

**MRI to Predict the Depth of Bowel Invasion**

Among the 40 patients, MRI suggested no bowel invasion (adherence only) in 15 cases and bowel invasion in 25 cases (12 cases with muscularis involvement and 13 cases with muscularis and submucosal  $\pm$  mucosal involvement).

Among the 25 patients with muscularis invasion on MRI, all had at least muscularis involvement at histopathology. Concerning the 13 patients with submucosal involvement on MRI, 9 (69%) had submucosal involvement at histopathology (Table 2).

The sensitivity of MRI for the detection of muscularis layer invasion was 68% (95% CI, 62%–68%), and the specificity was 100% (95% CI, 32%–100%). The PPV and NPV were 100% and 20%, respectively, and the negative LR was 0.32 (95% CI, 0.32%–1.17%) (Table 4).

For submucosal ( $\pm$  mucosal) layer invasion, the sensitivity of MRI was 47% (95% CI, 30%–61%), and the specificity was 81% (95% CI, 65%–93%). The PPV and NPV were 69% and 63%, respectively, and the positive and negative LRs were 2.49 (95% CI, 0.84%–8.57%) and 0.65 (95% CI, 0.42%–1.09%), respectively (Table 4). When considering the

**Table 1**

Baseline Patient Characteristics

Characteristics	
Median age, years (range)	33 (25–45)
Dysmenorrhea, n (%)	32 (80)
Dyspareunia, n (%)	27 (68)
Dyschezia, n (%)	28 (70)
Dysuria, n (%)	11 (28)
Chronic pelvic pain, n (%)	28 (70)
Infertility, n (%)	19 (48)

**Table 2**

Depth of Colorectal Infiltration of Endometriosis on Magnetic Resonance Imaging (MRI) and Histopathology					
Diagnosis on Histopathology					
MRI Diagnosis	No Digestive Infiltration, n	Muscularis, n	Muscularis + Submucosa, n	Muscularis + Submucosa + Mucosa, n	Total, n
No digestive infiltration	3	6	6	0	15
Muscularis	0	8	3	1	12
Muscularis + submucosa	0	0	2	0	2
Muscularis + submucosa + mucosa	0	4	7	0	11
Total	3	18	18	1	40

prediction of infiltration depth according to 3 categories (no lesion, muscularis, and submucosal/mucosal layers), the number of observed agreements was 20 (50% of the observations).

### **RES to Predict the Depth of Bowel Invasion**

RES diagnosed 14 patients with muscularis involvement; 23 patients with muscularis and submucosal involvement; and 3 patients with muscularis, submucosal, and mucosal involvement. Among the 14 patients with muscularis invasion identified by RES, 8 (57%) had muscularis involvement at histopathology, and 4 (29%) also had submucosal or mucosal involvement. Concerning the 26 patients with submucosal or mucosal impairment identified by RES, 15 (58%) had submucosal or mucosal involvement at histopathology (Table 3).

Concerning the specificity of RES for the detection of muscularis layer invasion, 3 cases were overestimated, but in those cases, invasion was described as very superficial adhesion of muscularis at RES. Typically, patients with suspected serosal endometriosis do not undergo segmental resection and should undergo serosal resection only and not digestive tract resection. In the current study, because of overestimation of bowel wall infiltration at the time of RES, these 3 patients underwent segmental resection when there was only serosal involvement or superficial invasion noted by pathology.

The sensitivity of RES for the detection of submucosal or mucosal invasion was 79% (95% CI, 61%–92%), and the

specificity was 48% (95% CI, 31%–60%). The PPV and NPV were 58% and 71%, respectively, and the positive and negative LRs were 1.51 (95% CI, 0.88%–2.24%) and 0.44 (95% CI, 0.13%–1.26%), respectively (Table 4). When considering the prediction of infiltration depth according to 3 categories (no lesion, muscularis, and submucosal/mucosal layers), the number of observed agreements was 23 (58% of the observations).

### **Combination of MRI and RES to Predict the Depth of Bowel Invasion**

When considering the prediction of infiltration depth according to 3 categories (no lesion, muscularis, and submucosal/mucosal layers), the number of observed agreements between RES and MRI was 12 (30% of the observations), and the intermethod agreement was  $-0.02$  (95% CI,  $-0.20\%$  to  $-0.15\%$ ).

MRI showed no bowel invasion (adherence only), whereas RES did in 15 patients (38%). On RES, one third (5/15) had muscularis layer involvement, and 10 had submucosal/mucosal layer involvement (Fig. 7). In this group of 5 patients with “no bowel invasion at MRI/muscularis layer involvement at RES,” histologic examination reported 1 patient with muscularis layer involvement, 2 patients with submucosal/mucosal involvement, and 2 patients without lesions (Fig. 8A and B).

Concerning the 10 patients without bowel invasion on MRI and submucosal/mucosal invasion on RES, histologic

**Table 3**

Depth of Colorectal Infiltration of Endometriosis on Rectal Endoscopic Sonography (RES) and Histopathology					
Diagnosis on Histopathology					
RES Diagnosis	No Digestive Infiltration, n	Muscularis, n	Muscularis + Submucosa, n	Muscularis + Submucosa + Mucosa, n	Total, n
Muscularis	2	8	4	0	14
Muscularis + submucosa	0	10	12	1	23
Muscularis + submucosa + mucosa	1	0	2	0	3
Total	3	18	18	1	40

**Table 4**

Magnetic Resonance Imaging (MRI) and Rectal Endoscopic Sonography (RES) Results

	MRI		RES	
	Muscularis	Submucosa/Mucosa	Muscularis	Submucosa/Mucosa
Sensitivity, %	68	47	—	79
Specificity, %	100	81	—	48
Positive predictive value, %	100	69	93	58
Negative predictive value, %	20	63	—	71
Positive likelihood ratio	—	2.49	—	1.51
Negative likelihood ratio	0.32	0.65	—	0.44

examination showed 5 patients with muscularis involvement, 4 patients with submucosal/mucosal involvement, and 1 patient without lesions. Finally, among the 15 patients without bowel invasion at MRI but with bowel invasion on RES, in 12 patients, histologic examination confirmed the diagnosis of bowel invasion.

**Discussion**

In the present study, we investigated whether MRI and/or RES could correctly determine patients who would benefit from rectal shaving because the procedure does not invade the colorectal submucosa. The PPV for MRI was superior to that of RES (100% vs 93%) for muscularis as well as for submucosal/mucosal invasion (69% vs 58%). RES showed higher sensitivity than MRI (79% vs 47%, respectively) for the detection of submucosal/mucosal infiltration. Based on the results of the present study, when MRI detects bowel involvement (either muscularis or submucosal/mucosal), RES is not required to confirm bowel infiltration. By contrast, when

not detected by MRI, RES is necessary to exclude bowel infiltration, particularly in patients showing symptoms of bowel DIE.

From a surgical point of view, the aim of surgical treatment of DIE could be to resect the entire lesion, which is associated with lower recurrence and less morbidity. Meuleman et al [4] reviewed the clinical outcome of surgical treatment of DIE in 49 studies and reported a low rate of recurrence with higher complications in the group that underwent bowel resection anastomosis (5.8% vs 17.6%) than the mixed surgical group (full-thickness resection and/or shaving). Studies have shown that rectal shaving was responsible for endometriosis lesion persistence in 84% of cases [18] and discoid resection in 42% of cases [19] compared with 31% of patients who underwent colorectal segmental resection [20].


The current study has several limitations. It is a retrospective study, and the classic bias encountered with such methodologies must be acknowledged. However, the radiologist who reinterpreted all MRIs and the sonographer who performed RES were both blinded to the results of histopathology as well as other imaging. Only 1 radiologist read all MRIs, and RES was performed by 1 sonographer. Another limitation is that the number of patients in the present study was relatively small, but the cohort of patients was homogeneous, and patients were managed consecutively, allowing a more reliable comparison. Only patients with a muscular lesion at least on RES were included, so it was not relevant to calculate the sensitivity, specificity, NPV, and LRs for the muscularis layer.

The final limitation of the study was that suboptimal MRI protocols were used in accordance with the recent European Society of Urogenital Radiology guidelines [21]; no systematic bowel preparation or antiperistaltic drugs were used. Gadolinium contrast was not standard protocol. For some radiologists, injection of gadolinium can be necessary to increase accuracy when there is ambiguity about a rectal lesion. With gadolinium, a 3-dimensional acquisition is used, allowing better spatial resolution and high-contrast resolution.

Regarding the 3 patients without bowel invasion (only adhesion) at histologic examination, negative histologic results have been reported by several authors [22,23] with rarefaction of glandular and stromal structures; thus, these results


**Fig. 7**

The depth of endometriotic involvement in the rectosigmoid on RES and MRI and on histopathology. SM-M = submucosa-mucosa.



**Fig. 8**

(A) MRI of the sagittal sequence T2-weighted, stenosis of the rectosigmoid junction. (B) Sagittal sequence T1-weighted showing the nodular lesion of the rectosigmoid explained by the presence of stools with hypointense filling the rectal ampulla. *Small arrow* = masked by the hypointense signal of the stools in the rectal ampulla.



should not refute the diagnosis of DIE. Interestingly, in the present study, the majority of discordant cases between MRI and histologic examination were related to underdiagnosis of the lesions (ie, depth of bowel invasion on histologic analysis was deeper than suspected on MRI). Such cases can be related to insufficient digestive preparation as shown in Fig. 8. This finding emphasizes the importance of digestive preparation by enema and rectal opacification to improve DIE infiltration depth assessment.

Our results regarding the capacity of RES to predict DIE colorectal infiltration depth are in accordance with previous studies [11,12,14,15]. Doniec et al [14] reported a sensitivity of 76% and 66% for the muscularis layer and the submucosal layer, respectively, with a disagreement rate of 25% in a series of 32 patients with colorectal endometriosis. Bazot et al [11] reported an agreement rate of 68% in a series of 54 cases of colorectal resection. Rossi et al [12] reported an agreement rate of 61% in a series of 38 cases, and

a rate of 56% was reported in the study from Roman et al (16 cases) [15] (Table 5).

In the past decade, MRI has been validated for the assessment of the number of lesions, location, size, and subsequent surgical resection [24–28]. Only 1 other study analyzed the capacity of MRI to predict the invasion depth of endometriosis on bowel wall infiltration. Busard et al [13] reported a sensitivity, specificity, PPV, and NPV of 100%, 75%, 96%, and 100%, respectively, for the detection of colorectal endometriosis lesions of the muscularis layer. Among the 22 patients with muscularis impairment on MRI, 11 (50%) had muscularis involvement only, and 9 (41%) had associated submucosal or mucosal involvement. Among the remaining 2 cases, 1 had serosal involvement, and the other had mucosal involvement [13].

Various methods have been studied to improve MRI accuracy, such as rectal or vaginal preparation, but without proven benefits [29,30]. Positive results were found with

**Table 5**

## Comparison of Rectal Endoscopic Sonography (RES) Studies

Study	Authors	N	Type of Study	Objective	Results
1	Bazot et al, 2007 [11]	81 (54 rectal DIE and 47 resections)	Cohort	RES and TVS for rectal DIE infiltration	Agreement 68%
2	Rossi et al, 2014 [12]	38	Retrospective, single center	RES for rectal DIE infiltration	Agreement 61%
3	Doniec et al, 2003 [20]	85 (32 rectal DIE and 25 resections)	Cohort	RES for rectal DIE infiltration	Disagreement 25% Sensitivity • Muscularis: 76% • Submucosa: 66%
4	Roman et al, 2008 [21]	16 (14 resections)	Retrospective single center	RES for rectal DIE infiltration	Agreement 56%

DIE = deep infiltrating endometriosis; TVS = transvaginal sonography.

Table 6

Comparison of Magnetic Resonance Imaging (MRI) Studies					
Study	Authors	N	Type of Study	Objective	Results
1	Busard et al, 2012 [13]	28	Retrospective, single center	MRI for rectal DIE infiltration depth	Sensitivity: 100% Specificity: 75% PPV: 96% NPV: 100%
2	Hottat et al, 2009 [27]	41	Prospective	3 Tesla MRI	Sensitivity: 96.3% Specificity: 100%
3	Manganaro et al, 2012 [28]	46	Prospective	3 Tesla MRI	Sensitivity: 93% Specificity: 75%

DIE = deep infiltrating endometriosis; NPV = negative predictive value; PPV = positive predictive value.

3-dimensional MRI, 3 Tesla MRI, and jelly methods [13,31,32]. Hottat et al [31] analyzed the contribution of 3 Tesla pelvic MRI in preoperative assessment and showed improved accuracy with a sensitivity and specificity of 96.3% and 100%, respectively, for the diagnosis of DIE (Table 6). Manganaro et al [32] reported the accuracy of 3 Tesla MRI in the evaluation of posterior cul-de-sac obliteration and showed a sensitivity and specificity of 93% and 75%, respectively. However, none of these new imaging modalities have been evaluated to predict the depth of bowel invasion.

## Conclusion

MRI is valuable for detecting colorectal DIE but is less accurate than RES in detecting submucosal/mucosal layer involvement and therefore cannot be used to determine whether patients should undergo segmental resection compared with a more conservative approach. When MRI detects bowel involvement (either muscularis or submucosal/mucosal layers), RES is not essential; however, if no intestinal lesion is detected by MRI, RES is necessary to confirm the diagnosis in symptomatic patients. Further large, prospective studies are necessary to determine whether MRI is sufficient for detecting the infiltration depth of colorectal endometriosis and ultimately the type of surgical repair best suited for this patient population.

## References

- Remorgida V, Ferrero S, Fulcheri E, Ragni N, Martin DC. Bowel endometriosis: presentation, diagnosis, and treatment. *Obstet Gynecol Surv*. 2007;62:461–470.
- Roman H, Ness J, Suciú N, et al. Are digestive symptoms in women presenting with pelvic endometriosis specific to lesion localizations? A preliminary prospective study. *Hum Reprod*. 2012;27:3440–3449.
- Darai E, Touboul C, Chéreau E, Bazot M, Ballester M. [Segmental resection for colorectal endometriosis: are there alternatives?]. *Gynecol Obstet Fertil*. 2012;40:116–120.
- Meuleman C, Tomassetti C, D'Hoore A, et al. Surgical treatment of deeply infiltrating endometriosis with colorectal involvement. *Hum Reprod Update*. 2011;17:311–326.
- De Cicco C, Corona R, Schonman R, Mailova K, Ussia A, Koninckx P. Bowel resection for deep endometriosis: a systematic review. *BJOG*. 2011;118:285–291.
- Donnez J, Squifflet J. Complications, pregnancy and recurrence in a prospective series of 500 patients operated on by the shaving technique for deep rectovaginal endometriotic nodules. *Hum Reprod*. 2010;25:1949–1958.
- Roman H, Vassilief M, Gourcerol G, et al. Surgical management of deep infiltrating endometriosis of the rectum: pleading for a symptom-guided approach. *Hum Reprod*. 2011;26:274–281.
- Roseau G, Dumontier I, Palazzo L, et al. Rectosigmoid endometriosis: endoscopic ultrasound features and clinical implications. *Endoscopy*. 2000;32:525–530.
- Bazot M, Thomassin I, Hourani R, Cortez A, Darai E. Diagnostic accuracy of transvaginal sonography for deep pelvic endometriosis. *Ultrasound Obstet Gynecol*. 2004;24:180–185.
- Hudelist G, Tuttlies F, Rauter G, Pucher S, Keckstein J. Can transvaginal sonography predict infiltration depth in patients with deep infiltrating endometriosis of the rectum? *Hum Reprod*. 2009;24:1012–1017.
- Bazot M, Malzy P, Cortez A, Roseau G, Amouyal P, Darai E. Accuracy of transvaginal sonography and rectal endoscopic sonography in the diagnosis of deep infiltrating endometriosis. *Ultrasound Obstet Gynecol*. 2007;30:994–1001.
- Rossi L, Palazzo L, Yazbeck C, et al. Can rectal endoscopic sonography be used to predict infiltration depth in patients with deep infiltrating endometriosis of the rectum? *Ultrasound Obstet Gynecol*. 2014;43:322–327.
- Busard MP, van der Houwen LE, Bleeker MC, et al. Deep infiltrating endometriosis of the bowel: MR imaging as a method to predict muscular invasion. *Abdom Imaging*. 2012;37:549–557.
- Doniec JM, Kahlke V, Peetz F, et al. Rectal endometriosis: high sensitivity and specificity of endorectal ultrasound with an impact for the operative management. *Dis Colon Rectum*. 2003;46:1667–1673.
- Roman H, Koureich K, Gromez A, Hochain P, Resch B, Marpeau L. Endorectal ultrasound accuracy in the diagnosis of rectal endometriosis infiltration depth. *Fertil Steril*. 2008;90:1008–1013.
- Bazot M, Nassar J, Darai E, et al. [Value of sonography and MR imaging for the evaluation of deep pelvic endometriosis]. *J Radiol*. 2005;86:461–467.
- Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33:159–174.
- Roman H, Opris I, Resch B, Tuech JJ, Sabourin JC, Marpeau L. Histopathologic features of endometriotic rectal nodules and the implications for management by rectal nodule excision. *Fertil Steril*. 2009;92:1250–1252.
- Remorgida V, Ragni N, Ferrero S, Anserini P, Torelli P, Fulcheri E. How complete is full thickness disc resection of bowel endometriotic

- lesions? A prospective surgical and histological study. *Hum Reprod.* 2005;20:2317–2320.
20. Badescu A, Roman H, Aziz M, et al. Mapping of bowel occult microscopic endometriosis implants surrounding deep endometriosis nodules infiltrating the bowel. *Fertil Steril.* 2016;105:430–434, e26.
  21. Bazot M, Bharwani N, Huchon C, et al. European Society of Urogenital Radiology (ESUR) guidelines: MR imaging of pelvic endometriosis. *Eur Radiol.* 2017;27:2765–2775.
  22. Bonte H, Chapron C, Vieira M, et al. Histologic appearance of endometriosis infiltrating uterosacral ligaments in women with painful symptoms. *J Am Assoc Gynecol Laparosc.* 2002;9:519–524.
  23. Albee RB, Sinervo K, Fisher DT. Laparoscopic excision of lesions suggestive of endometriosis or otherwise atypical in appearance: relationship between visual findings and final histologic diagnosis. *J Minim Invasive Gynecol.* 2008;15:32–37.
  24. Bazot M, Darai E, Hourani R, et al. Deep pelvic endometriosis: MR imaging for diagnosis and prediction of extension of disease. *Radiology.* 2004;232:379–389.
  25. Bazot M, Lafont C, Rouzier R, Roseau G, Thomassin-Naggara I, Darai E. Diagnostic accuracy of physical examination, transvaginal sonography, rectal endoscopic sonography, and magnetic resonance imaging to diagnose deep infiltrating endometriosis. *Fertil Steril.* 2009;92:1825–1833.
  26. Valentini AL, Gui B, Miccò M, et al. How to improve MRI accuracy in detecting deep infiltrating colorectal endometriosis: MRI findings vs. laparoscopy and histopathology. *Radiol Med.* 2014;119:291–297.
  27. Schneider C, Oehmke F, Tinneberg H-R, Krombach GA. MRI technique for the preoperative evaluation of deep infiltrating endometriosis: current status and protocol recommendation. *Clin Radiol.* 2016;71:179–194.
  28. Thomassin I, Bazot M, Detchev R, Barranger E, Cortez A, Darai E. Symptoms before and after surgical removal of colorectal endometriosis that are assessed by magnetic resonance imaging and rectal endoscopic sonography. *Am J Obstet Gynecol.* 2004;190:1264–1271.
  29. Bazot M, Gasner A, Lafont C, Ballester M, Darai E. Deep pelvic endometriosis: limited additional diagnostic value of postcontrast in comparison with conventional MR images. *Eur J Radiol.* 2011;80:e331–e339.
  30. Uyttenhove F, Langlois C, Collinet P, et al. Deep infiltrating endometriosis: should rectal and vaginal opacification be systematically used in MR imaging? *Gynecol Obstet Fertil.* 2016;44:322–328.
  31. Hottat N, Larrousse C, Anaf V, et al. Endometriosis: contribution of 3.0-T pelvic MR imaging in preoperative assessment—initial results. *Radiology.* 2009;253:126–134.
  32. Manganaro L, Vittori G, Vinci V, et al. Beyond laparoscopy: 3-T magnetic resonance imaging in the evaluation of posterior cul-de-sac obliteration. *Magn Reson Imaging.* 2012;30:1432–1438.