

Bibliografía

- Ismail S, Udd M, Lindström O, Rainio M, Halttunen J, Kylänpää L. Criteria for difficult biliary cannulation: start to count. *Eur J Gastroenterol Hepatol*. 2019 Oct; 31(10): 1200–5.
- Williams EJ, Taylor S, Fairclough P, Hamlyn A, Logan RF, Martin D, et al. Are we meeting the standards set for endoscopy? Results of a large-scale prospective survey of endoscopic retrograde cholangio-pancreatograph practice. *Gut*. 2007 Jun; 56(6): 821–9.
- Dumonceau JM, Kapral C, Aabakken L, et al. ERCP-related adverse events: European Society of Gastrointestinal Endoscopy (ESGE) guideline. *Endoscopy*. 2020 Feb; 52(2): 127–49.
- Mammen A, Haber G. Difficult biliary access: advanced cannulation and sphincterotomy technique. *Gastrointest Endosc Clin N Am*. 2015 Oct; 25(4): 619–30.
- Sasahira N, Kawakami H, Isayama H, et al. Early use of doubleguidewire technique to facilitate selective bile duct cannulation: the multicenter randomized controlled EDUCATION trial. *Endoscopy*. 2015 May; 47(5): 421–9.
- Tse F, Yuan Y, Moayyedi P, Leontiadis GI, Barkun AN. Double-guidewire technique in difficult biliary cannulation for the prevention of post-ERCP pancreatitis: a systematic review and meta-analysis. *Endoscopy*. 2017 Jan; 49(1): 15–26.
- Ito K, Fujita N, Noda Y, et al. Can pancreatic duct stenting prevent post-ERCP pancreatitis in patients who undergo pancreatic duct guidewire placement for achieving selective biliary cannulation? A prospective randomized controlled trial. *J Gastroenterol*. 2010 Nov; 45(11): 1183–91

Bibliografía

- Goff JS. Common bile duct pre-cut sphincterotomy: transpancreatic sphincter approach. *Gastrointest Endosc.* 1995 May; 41(5): 502–5.
- Sugiyama H, Tsuyuguchi T, Sakai Y, Mikata et al. Transpancreatic precut papillotomy versus double-guidewire technique in difficult biliary cannulation: prospective randomized study. *Endoscopy.* 2018 Jan; 50(1): 33–9.
- Goff JS. Common bile duct pre-cut sphincterotomy: transpancreatic sphincter approach. *Gastrointest Endosc.* 1995 May; 41(5): 502–5.
- Sakai Y, Tsuyuguchi T, Sugiyama H, et al. Transpancreatic precut papillotomy in patients with difficulty in selective biliary cannulation. *Hepatogastroenterology.* 2011 Nov–Dec; 58(112): 1853–8.
- Pécsi D, Farkas N, Hegyi P, et al. Transpancreatic sphincterotomy has a higher cannulation success rate than needle-knife precut papillotomy meta-analysis. *Endoscopy.* 2017 Sep; 49(9): 874–87.
- Liao WC, Angsuwatcharakon P, Isayama H, et al. International consensus recommendations for difficult biliary access. *Gastrointest Endosc.* 2017 Feb; 85(2): 295–304.
- Jang SI, Kim DU, Cho JH, Jeong S, Park JS, Lee DH, et al. Primary needle-knife fistulotomy versus conventional cannulation method in a high-risk cohort of post-endoscopic retrograde cholangiopancreatography pancreatitis. *Am J Gastroenterol.* 2020 Apr; 115(4): 616–24.

Bibliografía

- Nennstiel S, Weber A, Frick G, et al. Drainage-related complications in percutaneous transhepatic biliary drainage: an analysis over 10 years. *J Clin Gastroenterol*. 2015 Oct; 49(9): 764–70.
- Hamada T, Yasunaga H, Nakai Y, et al. Severe bleeding after percutaneous transhepatic drainage of the biliary system: effect of antithrombotic agents: analysis of 34,606 cases from a Japanese nationwide administrative database. *Radiology*. 2015 Feb; 274(2): 605–13
- Wang K, Zhu J, Xing L, Wang Y, Jin Z, Li Z. Assessment of efficacy and safety of EUS-guided biliary drainage: a systematic review. *Gastrointest Endosc*. 2016 Jun; 83(6): 1218–27.
- Sharaiha RZ, Khan MA, Kamal F, et al. Efficacy and safety of EUS-guided biliary drainage in comparison with percutaneous biliary drainage when ERCP fails: a systematic review and metaanalysis. *Gastrointest Endosc*. 2017 May; 85(5): 904–14.
- Dhir V, Itoi T, Khashab MA, et al. Multicenter comparative evaluation of endoscopic placement of expandable metal stents for malignant distal common bile duct obstruction by ERCP or EUS-guided approach. *Gastrointest Endosc*. 2015 Apr; 81(4): 913–23.